



## DESCRIPTION

### DRUG FOR ALLEVIATING MIGRAINE

#### TECHNICAL FIELD

5           The present invention relates to an external drug for dermal application, such as ointments or patches, in more detail, patches comprising in mixing l-menthol and an essential oil into a base containing a hydrophilic high-molecular weight compound, a polyhydric alcohol and water,  
10       which have a migraine-alleviating effect.

#### BACKGROUND ART

          The cause of migraine is not clear. It is suspected that increased blood stream due to expansion of head or  
15       cervix blood vessels caused by hormone unbalance, and muscle contraction in the area is the cause of migraine.

          For the treatment of migraine, analgesics for an internal application, which contain ergotamine tartrate, dimethothiazine mesylate, caffeine, etc., as an active  
20       ingredient are used. However, such a drug is often administered for long term and therefore, there is a possibility to induce side effects such as, anaphylaxis, insomnia, or gastrointestinal disorder.

          Accordingly, various preparations for dermal  
25       applications for treating migraine have been worked out.

For example, in Japanese Patent Publication B 6-67835,  
a composition where methysergide, an anti-serotonin, is  
dispersed in a hydrophilic polymer for systemic dermal  
application to prevent migraine is disclosed. Furthermore,  
5 in Japanese Patent Publication A Tokuhyo Hei 8-509749, a  
dermally therapeutic system containing sumatriptan, which  
is useful for migraine, cluster headache, etc., is  
disclosed.

However, these preparations for dermal application  
10 have a possibility to induce side effects, such as skin  
irritation, etc., by administering them for long term and  
therefore, these preparations are not favorable.

In addition, it is known that essential oils  
alleviate headache when used as an aromatherapy, but they  
15 are not simple to use.

The present inventors have extensively studied in  
order to obviate the above mentioned demerits, and as a  
result, have unexpectedly found that migraine can be  
alleviated by dermally administering to a human a drug  
20 containing l-menthol and an essential oil as active  
ingredients. Thus, the present invention has been  
completed.

#### DISCLOSURE OF INVENTION

25 The drug having a migraine-alleviating effect of the

present invention is a drug for a locally dermal application containing l-menthol and an essential oil as active ingredients. Its preferable preparations are ointments or patches, especially patches comprising l-menthol and an essential oil as active ingredients in a base containing hydrophilic high-molecular weight compound, a polyhydric alcohol and water.

The drug of the present invention is prepared by mixing l-menthol and an essential oil with a known base and if necessary, surfactants, preservatives, etc. to make ointments or patches by the conventional method.

The amount of l-menthol is for example, 0.01% - 1% by weight per total weight of base, preferably 0.05% - 0.5% by weight per total weight of base.

The essential oils used in the present invention are lavender oil, juniper oil, peppermint oil, rose oil, rosemary oil, etc. or a mixture thereof. The amount of these oils is 0.001% - 1% by weight per total weight of base, preferably 0.005 - 0.5% by weight per total weight of base.

In ointments, known bases such as white vaseline, yellow vaseline, lanolin, purified beeswax, cetanol, stearyl alcohol, hydrogenated oil, hydrocarbon gel, polyethylene glycol, etc. are used. To these bases, l-menthol and an essential oil and if necessary, surfactants,

preservatives, purified water, etc. are mixed to prepare ointments.

The especially preferable preparations of the present invention are patch-preparations which are prepared by  
5 mixing l-menthol and an essential oil as active ingredients into the base containing a hydrophilic high-molecular weight compound, polyhydric alcohol and water.

The patches of the present invention are in more detail explained as follows.

10 The hydrophilic high-molecular weight compounds used in the patches include, for example, gelatin, polyacrylic acid and its salt, polyvinyl alcohol, polyvinylpyrrolidone, carboxyvinyl polymer, sodium carboxymethyl cellulose, hydroxypropyl cellulose, methyl cellulose, ethyl cellulose,  
15 methyl vinyl ether-maleic acid anhydride copolymer, sodium alginate, poly ethylene oxide, acacia gum, xanthan gum, tragacanth gum, etc. These may be used in a mixture thereof.

The amount of the hydrophilic high-molecular weight  
20 compound is not limited, but when its amount is less than 2% by weight per total weight of base, the base is lack in viscosity not to become paste. On the other hand, when its amount is more than 20% by weight per total weight of base, it may occur that viscosity of the base becomes too  
25 high to smoothly prepare the preparation. Therefore, the

amount of the hydrophilic high-molecular weight compound is 2-20% by weight per total weight of base, preferably 5-15% by weight per total weight of base.

5 The polyhydric alcohols include glycerin, sorbitol, propylene glycol, polyethylene glycol, 1,3-butylene glycol, ethylene glycol, etc. These may be used in a mixture thereof.

The amount of the polyhydric alcohol is 8 - 60% by weight per total weight of base, preferably 10 - 50% by weight per total weight of base.

10 When its amount is less than 8% by weight per total weight of base, humidity-keeping effect becomes poor and water become volatile in short times. On the other hand, when its amount is more than 60% by weight per total weight of base, it is difficult to mix with other substances and to use the polyhydric alcohol so much is not desirable.

15 The amount of water is 20 - 80% by weight per total weight of base, preferably 25 - 70% by weight per total weight of base.

20 When its amount is less than 20% by weight per total weight of base, dissolution of the hydrophilic high-molecular weight compound is not satisfactory and it is impossible to homogeneously extend the base. On the other hand, when its amount is more than 80% by weight per total

weight of base, it may occur that the base becomes too soft to spread out. Therefore, it is not desirable to use water so much.

5 The amount of l-menthol is 0.01-1% by weight per total weight of base, preferably 0.05-0.5% by weight per total weight of base as mentioned above. The amount of the essential oil is 0.001-1% by weight per total weight of base, preferably 0.005-0.5% by weight per total weight of base as mentioned above.

10 In addition to the above mentioned base, following additives which are usually used in patches can be mixed in the usual amount: excipients (kaolin, bentonite, titanium oxide, etc.), surfactants (glycerin fatty acid ester, polyoxyethylene castor oil, polyoxyethylene  
15 hydrogenated castor oil, sorbitan fatty acid ester, polysorbate 80, polysorbate 60, sorbitan sesquioleate), crosslinking agents (multivalent metal such as aluminum hydroxide, aluminum glycinate, dihydroxyaluminum aminoacetate, synthetic hydrotalcite, etc.), coloring  
20 agents (new coccin, tartrazine, brilliant blue FCF), and preservatives (p-hydroxybenzoic acid ester, sorbic acid salt, isopropyl methyl phenol, hinokitiol, phenoxyethanol, etc.)

25 The base is prepared by mixing each ingredient in accordance with the conventional method. For example, a

part of a hydrophilic high-molecular weight compound and a polyhydric alcohol are dissolved in water, and if desired, other additives are mixed, and then l-menthol and an essential oil are added to the mixture to be kneaded.

5 Then, residual of the hydrophilic high-molecular weight compound and other additives are mixed thereto to prepare the base.

The base thus prepared is spread on an appropriate support and a releasing paper is put on the base in order  
10 to protect the base. The base cut in a fixed size to prepare desired patches.

The amount of the base in patches is  $200-5000\text{g/m}^2$ , preferably  $500-2000\text{g/m}^2$ .

The support is one such as non-woven fabrics, fabrics,  
15 knits, etc., used in usual patches. Its material is an synthetic fiber such as nylon, rayon, polyester, polypropylene, etc. or a natural fiber such as cotton. As the releasing paper, plastic film such as polyethylene film, etc. and others used in usual patches are used.

20 Shapes of the patches may be ellipse, rectangle, triangle, boomerang type, facemask type, etc.

The patches of the present invention are preferably applied to on forehead, nape of the neck, temple, a half of face and/or full face, and by doing the patch thereto,  
25 migraine-alleviating effect effectively appears.

# BEST MODE FOR CARRYING OUT THE INVENTION

The present invention and its effect are illustratively explained by working examples and tests, but the invention should not be limited by these examples.

## Examples 1-6

Using ingredients shown in Tables 1 and 2, patches (Examples 1-6) were prepared by the conventional method. Namely, a part of the hydrophilic high-molecular weight compound and the polyhydric alcohol were dissolved in purified water, and if necessary, other ingredients were added thereto. The mixture was fully kneaded. Then, l-menthol and the essential oil were added to the mixture and further, the residue of the hydrophilic high-molecular weight compound and other ingredients were added to. Finally, the residue of the purified water was added to the mixture. The mixture was homogeneously kneaded to prepare a base.

The base prepared was spread on the support (1000g/m<sup>2</sup>) and a releasing paper or plastic film was put on it. The base was cut into a fixed size to prepare patches.

The bases prepared above, as such may be used as ointments.



Table 1.

Ingredients	Percent by Weight		
	Example 1	Example 2	Example 3
Polyacrylic acid	1.0	2.5	1.25
Sodium polyacrylate	5.0	6.0	6.0
Sodium carboxy methylcellulose	5.0	4.0	5.5
Gelatin	0.4	-	0.2
Polyvinyl alcohol	0.2	-	-
Tartaric acid	0.2	0.15	0.25
Disodium edetate	0.1	0.08	0.07
Glycerin	22.0	15.0	18.0
70% Sorbitol solution	-	15.0	-
Aluminum hydroxide	0.3	-	-
Synthetic hydrotalcite	-	0.2	-
Dihydroxyaluminum acetate	-	-	0.08
Polysorbate 80	0.1	0.1	0.1
Caster oil	0.5	0.5	0.5
Methylparaben	0.1	0.1	0.1
l-Menthol	0.3	0.15	0.1
Peppermint oil	0.2	-	-
Rose oil	-	0.1	-
Lavender oil	-	-	0.01
Purified water	Residue	Residue	Residue
	100	100	100

Table 2.

Ingredients	Percent by Weight		
	Example 4	Example 5	Example 6
Polyacrylic acid	1.5	2.0	1.25
Sodium polyacrylate	5.0	5.5	6.0
Sodium carboxy methylcellulose	5.0	4.0	5.5
Gelatin	-	-	-
Polyvinyl alcohol	0.2	-	-
Tartaric acid	0.2	0.15	0.3
Disodium edetate	0.1	0.08	0.07
Glycerin	20.0	15.0	20.0
70% Sorbitol solution	10.0	15.0	-
Aluminum hydroxide	0.3	-	-
Synthetic hydrotalcite	0.15	-	-
Dihydroxyaluminum acetate	-	0.1	0.1
Polysorbate 80	0.1	0.1	0.1
Caster oil	0.5	0.5	0.5
Methylparaben	0.1	0.1	0.1
l-Menthol	0.8	0.25	0.05
Peppermint oil	0.2	0.4	-
Rose oil	-	0.4	0.05
Lavender oil	0.05	-	0.1
Purified water	Residue	Residue	Residue
	100	100	100

Comparative example 1

The patch was prepared by the same method as Example  
5 1 using the same ingredients as Example 1 provided that

the same amount of water as l-menthol was used instead of l-menthol (Only an essential oil is used as an active ingredient).

5 Comparative example 2

The patch was prepared by the same method as Example 1 using the same ingredients as Example 1 provided that the same amount of water as an essential oil was used instead of the essential oil (Only l-menthol is used as an  
10 active ingredient).

Comparative example 3

The patch was prepared by the same method as Example 1 using the same ingredients as Example 1 provided that  
15 the same amount of water was used instead of the essential oil and l-menthol (Any active ingredient was not used).

Next, each two patches (5x7cm) of Examples 1, 3, 5 and Comparative examples 1-3 were put on each volunteer.  
20 The following items were sensitively evaluated.

Test 1

On their foreheads of ten volunteers suffering from migraine were put each patch of Examples 1, 3, 5 and  
25 Comparative examples 1-3, and migraine-alleviating effect

was evaluated by sensory test under following evaluation-standards.

Evaluation-standard on effects

Point 1: no effect

5 Point 2: weak effect

Point 3: effective

Point 4: clearly effective

Point 5: strongly effective

10 Efficacy (point) was indicated by the average of volunteer's evaluations. The duration of the effect was indicated by the average of volunteer's reported times.

The result was shown in the following Table 3.

Table 3

	Efficacy(point)	Duration of effect(hour)
Example 1	4.2	7.3
Example 3	4.3	7.9
Example 5	3.9	6.5
Comparative example 1	2.5	3.2
Comparative example 2	2.8	2.8
Comparative example 3	1.3	2.1

15 As is clear from the result of Table 3, patches of Examples 1, 3 and 5 were superior in efficacy (point) to

patches of Comparative examples 1-3, and therefore, it is recognized that patches of Examples 1, 3 and 5 are excellent in migraine-alleviating effect and that its effect lasts for long hours.

5

## Test 2

On various regions of ten volunteers suffering from migraine were put each patch of Examples 1, 3, 5 and Comparative example 3, and migraine-alleviating effect depending on the region was evaluated by sensory test under following evaluation-standards.

Evaluation standard:

+ : Positive alleviating efficacy

± : Weak alleviating efficacy

15 - : No alleviating efficacy

The result is shown in Table 4.

Table 4.

Application region	Alleviating efficacy			
	Example 1	Example 3	Example 5	Comparative example 3
Forehead	+	+	+	±
Nape of neck	+	+	+	-
Temple	+	+	+	±
Shoulder	±	±	±	-
Back	-	-	-	-
Breast	-	-	-	-

As is clear from the result of Table 4, when applying the preparations of the present invention, that is preparations of Examples 1, 3 and 5 to face, nape of the neck and temple, the preparations were recognized being superior in migraine-alleviating efficacy. On the other hand, a patch of Comparative example 3 hardly showed migraine-alleviating efficacy in any region.

#### INDUSTRIAL APPLICABILITY

The preparation of the present invention is excellent in migraine-alleviating efficacy, and even when using for long terms, there is hardly a possibility to induce side effects and the preparation of the present invention is very convenient and useful.